

BECAUSE YOU CAN'T SEE STERILE.^{5M}

2014

www.CrosstexBMS.com

Contact us for test status at 800-819-3336

How often should I perform biological monitoring (BI) (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenence.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

		JA	NUA	RY		
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

FEBRUARY							
S	M	T	W	Т	F	S	
						1	
2	3	4	5	6	7	8	
9	10	11	12	13	14	15	
16	17	18	19	20	21	22	
23	24	25	26	27	28		

MARCH						
S	M	T	W	T	F	S
2	3	4	5	6	7	1 8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

TEST DATE	INITIALS OF PERSON WHO RAN TEST	MAIL DATE	INITIALS OF PERSON WHO MAILED TEST

Have any Questions?
Call a Crosstex Customer Care at:

1-800-819-3336



BECAUSE YOU CAN'T SEE STERILE. SM

2014

www.CrosstexBMS.com

Contact us for test status at 800-819-3336

How often should I perform biological monitoring (BI) (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenence.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

	APRIL					
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			

			MAY			
S	M	T	W	Т	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

			JUNE			
S	M	T	W	Т	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

TEST DATE	INITIALS OF PERSON WHO RAN TEST	MAIL DATE	INITIALS OF PERSON WHO MAILED TEST

Have any Questions?
Call a Crosstex Customer Care at:

1-800-819-3336



BECAUSE YOU CAN'T SEE STERILE.™

2014

www.CrosstexBMS.com

Contact us for test status at 800-819-3336

How often should I perform biological monitoring (BI) (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenence.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was

OCTOBER					periorilled	
S	M	T	W	Т	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

	NOVEMBER							
S	M	T	W	T	F	S		
						1		
2	3	4	5	6	7	8		
9	10	11	12	13	14	15		
16	17	18	19	20	21	22		
23	24	25	26	27	28	29		
30								

	DECEMBER						
S	M	T	W	Т	F	S	
	1	2	3	4	5	6	
7	8	9	10	11	12	13	
14	15	16	17	18	19	20	
21	22	23	24	25	26	27	
28	29	30	31				

INITIALS INITIALS **TEST DATE MAIL DATE** OF PERSON WHO RAN TEST OF PERSON WHO MAILED TEST

Have any Questions?
Call a Crosstex Customer Care at:

1-800-819-3336



BECAUSE YOU CAN'T SEE STERILE.^{5M}

2014

www.CrosstexBMS.com

Contact us for test status at 800-819-3336

How often should I perform biological monitoring (BI) (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenence.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

			JULY			performed
S	M	T	W	Т	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

AUGUST									
S	M	T	W	T	F	S			
					1	2			
3	4	5	6	7	8	9			
10	11	12	13	14	15	16			
17	18	19	20	21	22	23			
24	25	26	27	28	29	30			
31									

SEPTEMBER									
S	M	Т	W	Т	F	S			
	1	2	3	4	5	6			
7	8	9	10	11	12	13			
14	15	16	17	18	19	20			
21	22	23	24	25	26	27			
28	29	30							

INITIALS INITIALS TEST DATE MAIL DATE OF PERSON WHO RAN TEST OF PERSON WHO MAILED TEST

Have any Questions?
Call a Crosstex Customer Care at:
1-800-819-3336